



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The Coalville Coal Field, Utah. By CARROLL H. WEGEMANN. U.S. Geol. Survey, Bull. 581-E, 1915. Pp. 24, pls. 6.

The Coalville coal field lies about 30 miles northeast of Salt Lake City, in the valley of Weber River. High-grade sub-bituminous coal has been mined here for more than fifty years.

The report covers four townships. The rocks of the district include some 8,000 feet of shales and sandstones of Colorado and Montana (?) age, which are folded into a slightly overturned and pitching anticline, and are unconformably overlain by 1,000 feet or more of Wasatch conglomerate. Several transverse and nearly vertical faults of small displacement cut the gently dipping limb of the fold. Both the folding and the faulting took place chiefly in pre-Wasatch time, when considerable erosion was likewise accomplished; but weaker movement of both types appears to have followed the deposition of the Wasatch beds.

The one productive coal bed, known as the "Wasatch" bed, varies from 5 to 12 feet in thickness. This coal compares favorably in quality with several Wyoming coals. Coal occurs in thinner seams at two other horizons, 2,000 feet above and 850 feet below the "Wasatch" bed, respectively. All three horizons are in the Cretaceous system.

C. W. T.

Preliminary Report on the Clay and Shale Deposits of the Province of Quebec. By J. KEELE. Canada Dept. of Mines, Memoir 64. Ottawa, 1915. Pp. 280+iv, pls. XXXIV, figs. 13, map 1.

Describes the clay-bearing horizons and groups producing localities by the age of the clay produced. Particular emphasis is laid on the Pleistocene clays. A considerable portion of the memoir is devoted to the technologic aspects of the clay industries.

A. D. B.

"The Pebble Phosphates of Florida." By E. H. SELLARDS. *Florida Geol. Survey, Seventh Annual Report*, 1915, pp. 25-116.

In an earlier report this writer has discussed the origin of hard-rock phosphates, and this paper extends the study to land-pebble and river-pebble deposits. The land-pebble phosphates are found in the Bone Valley formation of late Miocene or early Pliocene age. They form a portion of a basal conglomerate laid down by a sea advancing over the Alum Bluff formation, a phosphatic marl of late Oligocene age. In this